

Teprotumumab-trbw (Tepezza) Patient Referral Form

Patient Demographics Information

Patient Name		SSN#	DOB
Patient Address			
Primary Phone	Cellular Phone	Work Phone	
Emergency Contact Name, Relationship		Emergency Contact Phone Number	

Additional Documentation Needed

- Copy of insurance cards
- Patient face sheet w/demographics
- History and Physical

Patient Insurance Information

Insurance Plan #1		Insurance Plan #2	
Plan Address		Plan Address	
Plan Phone & Fax Numbers		Plan Phone & Fax Numbers	
Subscriber Name	DOB	Subscriber Name	DOB
Policy Number	Group ID	Policy Number	Group ID

Patient Clinical Information

Gender <input type="checkbox"/> M <input type="checkbox"/> F	Height (inches)	Weight (lbs.)	Allergies (food/drug)
---	-----------------	---------------	-----------------------

Statement of Medical Necessity / Primary Diagnosis

ICD10:	Description of diagnosis:
--------	---------------------------

Medication Information / Prescription and Orders

Medication: Tepezza 500 mg	Directions: Infuse IV per manufacturer guidelines OR over _____ hours. Prior to infusion, reconstitute each Tepezza 500 mg vial with 10 mL of Sterile Water for Injection and mix in a NaCl 0.9% bag for total 100 mL for doses <1800 mg or 250 mL for doses ≥1800 mg
---	---

Initial dose <input type="checkbox"/> _____ mg (10 mg/kg) IV once #1 dose, 21-day supply, no refill	*If subsequent treatment cycles only Date of last infusion:
Subsequent doses 2-8 <input type="checkbox"/> _____ mg (20 mg/kg) IV every 3 weeks #1 dose, 21 day supply, refill x 6	Next dose due:

Line Access RN to start peripheral IV or use existing CVAD. RN to administer catheter flushing and medications as directed below. <input type="checkbox"/> Sodium Chloride 0.9% 10 mL Flush: Flush with 2-10 mL sodium chloride 0.9% IV before and after medication administration or every 24 hours while IV access in place. <input type="checkbox"/> Sodium Chloride 0.9% 10 mL STERILE Flush: Flush with 5-10 mL STERILE sodium chloride 0.9% IV as needed for port access. <input type="checkbox"/> Heparin Lock 100U/mL 5 mL Flush: Lock with 5 mL heparin 100U/mL after each use or daily while port accessed.	Quantity/Refills Dispense: quantity #QS + PRN refills unless otherwise noted <input type="checkbox"/> Other
--	---

Adverse Reaction Orders

*** If infusion reaction occurs pause infusion for 10 minutes, resume infusion at previously tolerated rate and use appropriate medical management. If reaction continues, stop infusion and notify physician.**

Prescriber Information

Physician Name		Office Contact	
Practice Address		Practice Phone	
NPI#	License #	DEA#	
Physician Signature Required - Substitution Permitted		Date	Physician Signature Required - Dispense as Written
			Date