

Rozanolixizumab-noli (Rystiggo) Patient Referral Form

Patient Demographics Information

Patient Name		SSN#	DOB
Patient Address			
Primary Phone	Cellular Phone	Work Phone	
Emergency Contact Name, Relationship		Emergency Contact Phone Number	

Additional Documentation Needed

- | | | |
|--|---|---|
| <input type="checkbox"/> Copy of insurance cards | <input type="checkbox"/> History and Physical | <input type="checkbox"/> CBC w/diff, BMP, & CMP |
| <input type="checkbox"/> Patient face sheet w/demographics | <input type="checkbox"/> Antibody testing results | <input type="checkbox"/> Recent vitals including blood pressure |

Patient Insurance Information

Insurance Plan #1		Insurance Plan #2	
Plan Address		Plan Address	
Plan Phone & Fax Numbers		Plan Phone & Fax Numbers	
Subscriber Name	DOB	Subscriber Name	DOB
Policy Number	Group ID	Policy Number	Group ID

Patient Clinical Information

Gender <input type="checkbox"/> M <input type="checkbox"/> F	Height (inches)	Weight (lbs.)	Allergies (food/drug)
---	-----------------	---------------	-----------------------

Statement of Medical Necessity / Primary Diagnosis

ICD10:	Description of diagnosis:	Current MG-ADL Score:
--------	---------------------------	-----------------------

Medication Information / Prescription and Orders

Medication: Rystiggo	Directions: Infuse SubQ with mechanical syringe pump at a rate of up to 20 mL/hr per manufacturer guidelines.	Quantity/Refills Dispense: 6-week supply, refill x12mo. Unless otherwise noted <input type="checkbox"/> Other
<input type="checkbox"/> < 50 kg <input type="checkbox"/> 420 mg (3 mL) SubQ every week for 6 weeks. <input type="checkbox"/> 50-99 kg <input type="checkbox"/> 560 mg (4 mL) SubQ every week for 6 weeks. <input type="checkbox"/> ≥100 kg <input type="checkbox"/> 840 mg (6 mL) SubQ every week for 6 weeks.		*If subsequent treatment cycles only Start date of last cycle:

Premedications

<input type="checkbox"/> Give premedication(s) 30 minutes prior to infusion (<i>generics will be dispensed</i>) Diphenhydramine <input type="checkbox"/> 25-50 mg PO Fexofenadine <input type="checkbox"/> 180 mg PO Acetaminophen <input type="checkbox"/> 325-650 mg PO OR <input type="checkbox"/> _____ mg PO <input type="checkbox"/> Lidocaine 2.5% and prilocaine 2.5% (EMLA) topical cream: apply to SQ needle site prior to access PRN pain upon needle insertion. <input type="checkbox"/> RN to instruct patient to hydrate pre/post infusion. <input type="checkbox"/> Other	Quantity/Refills Dispense: Quantity #QS + PRN refills unless otherwise noted <input type="checkbox"/> Other
---	---

Adverse Reaction Orders

In the event of an infusion reaction (i.e. fever, chills, backache, headache, rigors, etc.) the following order will be followed and the ordering provider will be notified:

- Mild reaction: Pause infusion and administer diphenhydramine 50 mg PO x1 dose. If needed, give an additional dose of diphenhydramine 50 mg PO x1 dose. Max 2 doses. Resume infusion once symptoms resolve.
- Moderate reaction: Stop infusion and administer diphenhydramine 50 mg PO x1 dose.
- Severe reaction (w/breathing problems): CALL 911, administer epinephrine 0.3 mg IM. #QS + PRN refills

Prescriber Information

Prescriber Name		Office Contact	
Practice Address		Practice Phone	
NPI#	License #	Practice Fax	
Physician Signature Required - Substitution Permitted		Date	Physician Signature Required - Dispense as Written
			Date