



**Patient Demographics Information**

Patient Name		SSN#	DOB
Patient Address			
Primary Phone	Cellular Phone	Work Phone	
Emergency Contact Name, Relationship		Emergency Contact Phone Number	

**Additional Documentation Needed**

<input type="checkbox"/> Copy of insurance cards	<input type="checkbox"/> History and Physical	<input type="checkbox"/> Recent vitals including blood pressure
<input type="checkbox"/> Patient face sheet w/demographics	<input type="checkbox"/> CBC w/diff, BMP, & CMP	<input type="checkbox"/> Confirmation of DMD gene mutation
<input type="checkbox"/> Kidney Function utilizing serum cystatin C, urine dipstick, or urine-to-creatinine ratio		

**Patient Insurance Information**

Insurance Plan #1		Insurance Plan #2	
Subscriber Name	DOB	Subscriber Name	DOB
Policy Number	Group ID	Policy Number	Group ID

**Patient Clinical Information**

Gender <input type="checkbox"/> M <input type="checkbox"/> F	Height (inches)	Weight (lbs.)	Allergies (food/drug)
---	-----------------	---------------	-----------------------

**Statement of Medical Necessity / Primary Diagnosis**

ICD10:	Description of diagnosis:
--------	---------------------------

**Medication Information / Prescription and Orders**

<b>Medication:</b>  Viltepso	<b>Directions:</b> Infuse intravenously per manufacturer guidelines over 60 minutes. *If calculated dose is less than 100 mL, dilute with sodium chloride 0.9% to a total volume of 100 mL. If calculated dose is 100 mL or more, further dilution not required.	<b>Quantity/Refills</b> Dispense: #4 doses (one month) + _____ refills <input type="checkbox"/> Other
<input type="checkbox"/> DMD dosing: _____ mg (80mg/kg) intravenously once weekly. *If patient is currently taking Viltepso, date of last dose: <input type="checkbox"/> Other dosing: _____ mg intravenously every ____ week(s)		

**Premedication(s)**

<input type="checkbox"/> Give premedication(s) 30 minutes prior to infusion (generics will be dispensed) diphenhydrAMINE <input type="checkbox"/> Pediatric Dosing: _____mg (1mg/kg/dose) PO <input type="checkbox"/> Adult Dosing: 25-50mg PO Acetaminophen <input type="checkbox"/> Pediatric Dosing: _____mg (10-15mg/kg/dose) PO <input type="checkbox"/> Adult Dosing: 325-650mg PO Other: <input type="checkbox"/> _____ (medication name): _____ (dosing)	<b>Quantity/Refills</b> Dispense: #QS + 12 refills unless otherwise noted <input type="checkbox"/> Other
<input type="checkbox"/> Lidocaine 2.5% and Prilocaine 2.5% topical cream: Apply to IV site prior to access PRN for pain upon needle insertion. <input type="checkbox"/> Other (prescriber to specify):	

**Line Access**

RN to start peripheral IV or use existing CVAD. RN to administer catheter flushing and medications per Heritage Biologics policy. <input type="checkbox"/> Sodium Chloride 0.9% 10 mL Flush: Flush with 2-10 mL sodium chloride 0.9% IV before and after medication administration or every 24 hours while IV access in place. <input type="checkbox"/> Sodium Chloride 0.9% 10 mL STERILE Flush: Flush with 5-10 mL STERILE sodium chloride 0.9% as needed for port access <input type="checkbox"/> Heparin Lock <b>10 units/mL</b> 5 mL Flush: Lock with 2-5 mL heparin <b>10 units/mL</b> after each use or daily while port accessed. <input type="checkbox"/> Heparin Lock <b>100 units/mL</b> 5 mL Flush: Lock with 2-5 mL heparin <b>100 units/mL</b> after each use or daily while port accessed.	<b>Quantity/Refills</b> Dispense: quantity #QS PRN refills unless otherwise noted <input type="checkbox"/> Other
---	---

**Adverse Reaction Orders**

If infusion reaction(s) occur, pause infusion for 10 minutes, then resume infusion at previously tolerated rate and use appropriate medical management  
If reaction(s) continue, stop infusion and notify physician.

**Prescriber Information**

Physician Name		Office Contact	
Practice Address			
NPI#	License #	Practice Phone	
Physician Signature Required - Substitution Permitted		Date	Physician Signature Required - Dispense as Written
			Date