



Patient Demographics Information

Patient Name		SSN#	DOB
Patient Address			
Primary Phone	Cellular Phone	Work Phone	
Emergency Contact Name, Relationship		Emergency Contact Phone Number	

Additional Documentation Needed

<input type="checkbox"/> Copy of insurance cards	<input type="checkbox"/> History and Physical	<input type="checkbox"/> Recent vitals including blood pressure
<input type="checkbox"/> Patient face sheet w/demographics	<input type="checkbox"/> CBC w/diff, BMP, & CMP	<input type="checkbox"/> Screening for HBV infection

Patient Insurance Information

Insurance Plan #1		Insurance Plan #2	
Subscriber Name	DOB	Subscriber Name	DOB
Policy Number	Group ID	Policy Number	Group ID

Patient Clinical Information

Gender <input type="checkbox"/> M <input type="checkbox"/> F	Height (inches)	Weight (lbs.)	Allergies (food/drug)
---	-----------------	---------------	-----------------------

Statement of Medical Necessity / Primary Diagnosis

ICD10:	Description of diagnosis:
--------	---------------------------

Medication Information / Prescription and Orders

Medication: Ocrevus Zunovo	Directions: Infuse SubQ per manufacturer guidelines over approximately 10 minutes.	Quantity/Refills Dispense: #1 dose + _____ refills <input type="checkbox"/> Other
Dosing: <input type="checkbox"/> MS dosing: 920 mg/23,00 units SubQ every 6 months	Date of Last Dose:	

Premedications

Give premedication(s) 30 minutes prior to infusion	Quantity/Refills Dispense: quantity #QS PRN refills (unless otherwise noted) <input type="checkbox"/> Other
Acetaminophen <input type="checkbox"/> 325-650 mg PO OR <input type="checkbox"/> _____ mg PO	
Dexamethasone <input type="checkbox"/> 20 mg PO	
Diphenhydramine <input type="checkbox"/> 25-50 mg PO	
Other <input type="checkbox"/> _____	

Adverse Reaction Orders

- Mild reaction: Pause infusion and resume after symptoms have resolved
- Moderate reaction: Pause infusion and give diphenhydrAMINE 50mg PO. Resume infusion after symptoms have resolved. Notify Pharmacist.
- Severe reaction (w/breathing problems): stop infusion, call 911 and administer EPHINEPHrine 0.3 mg or 0.15 mg IM (as determined by patient weight).

Prescriber Information

Physician Name		Office Contact	
Practice Address			
NPI#	License #	Practice Phone	
Physician Signature Required - Substitution Permitted	Date	Physician Signature Required - Dispense as Written	Date