



# Ocrelizumab (Ocrevus) Patient Referral Form

Admissions Fax # 844-878-6917

Admissions Phone # 855-WE-R-RARE (855-937-7273)

## Patient Demographics Information

|                                      |                |                                |     |
|--------------------------------------|----------------|--------------------------------|-----|
| Patient Name                         |                | SSN#                           | DOB |
| Patient Address                      |                |                                |     |
| Primary Phone                        | Cellular Phone | Work Phone                     |     |
| Emergency Contact Name, Relationship |                | Emergency Contact Phone Number |     |

## Additional Documentation Needed

- |                                     |                          |                                          |
|-------------------------------------|--------------------------|------------------------------------------|
| - Copy of insurance cards           | - History and Physical   | - Recent vitals including blood pressure |
| - Patient face sheet w/demographics | - CBC w/diff, BMP, & CMP | - Screening for HBV infection            |

## Patient Insurance Information

|                   |          |                   |          |
|-------------------|----------|-------------------|----------|
| Insurance Plan #1 |          | Insurance Plan #2 |          |
| Subscriber Name   | DOB      | Subscriber Name   | DOB      |
| Policy Number     | Group ID | Policy Number     | Group ID |

## Patient Clinical Information

|                                                                 |                 |              |                       |
|-----------------------------------------------------------------|-----------------|--------------|-----------------------|
| Gender<br><input type="checkbox"/> M <input type="checkbox"/> F | Height (inches) | Weight (lbs) | Allergies (food/drug) |
|-----------------------------------------------------------------|-----------------|--------------|-----------------------|

## Statement of Medical Necessity / Primary Diagnosis

|        |                           |
|--------|---------------------------|
| ICD10: | Description of diagnosis: |
|--------|---------------------------|

## Medication Information / Prescription and Orders

|                                                                                                                                                                                                                                                                                               |                                                                                                                                        |                                                                |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------|
| <b>Medication:</b><br>Ocrevus                                                                                                                                                                                                                                                                 | <b>Directions:</b><br>Infuse IV per manufacturer guidelines <b>OR</b> over _____ hours.<br>Titration rate according to package insert. | *If subsequent treatment cycles only<br>Date of last infusion: |
| <input type="checkbox"/> Initial treatment cycle<br>300 mg IV day 1, 14 day drug-free period, 300 mg IV day 15, #2 doses, refill 0<br><input type="checkbox"/> Subsequent treatment cycles<br>600 mg IV every 6 months, #1 dose, refill x12mos <b>OR</b> <input type="checkbox"/> Other _____ |                                                                                                                                        | Next dose due:                                                 |

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |                                                                                                                             |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|
| <b>Line Access</b><br>Line Type: <input type="checkbox"/> PIV <input type="checkbox"/> Port/CVAD<br>RN to start peripheral IV or use existing CVAD. RN to administer catheter flushing and medications per orders below.<br><input type="checkbox"/> Sodium Chloride 0.9% 10 mL Flush: Flush with 2-10 mL sodium chloride 0.9% IV before and after medication administration or every 24 hours while IV access in place.<br><input type="checkbox"/> Sodium Chloride 0.9% 10 mL STERILE Flush: Flush with 5-10 mL STERILE sodium chloride 0.9% IV as needed for port access.<br><input type="checkbox"/> Heparin Lock 100U/mL 5 mL Flush: Lock with 5 mL heparin 100U/mL after each use or daily while port accessed.                                                                                                                                  | <b>Quantity/Refills</b><br>Dispense:<br>Quantity #QS + PRN refills unless otherwise noted<br><input type="checkbox"/> Other |
| <b>Premedications</b><br><input type="checkbox"/> Give premedication 30 minutes prior to infusion ( <i>generics will be dispensed</i> )<br>Diphenhydramine: <input type="checkbox"/> 25-50 mg PO<br>Methylprednisolone: <input type="checkbox"/> 125 mg slow IV push over 5 minutes<br>Acetaminophen: <input type="checkbox"/> 325-650 mg PO <b>OR</b> <input type="checkbox"/> _____ mg PO<br><input type="checkbox"/> EMLA topical cream (Lidocaine 2.5% and Prilocaine 2.5%): apply to IV site prior to access PRN for pain upon needle insertion.<br><input type="checkbox"/> Sodium Chloride 0.9% 500 mL - 1000 mL IV over 1-2 hours as tolerated daily PRN for hydration.<br><input checked="" type="checkbox"/> RN to instruct patient to hydrate pre/post infusion.<br><input type="checkbox"/> Other (Physician to specify):                  |                                                                                                                             |
| <b>Adverse Reaction Orders</b><br>In the event of an infusion reaction (ie: fever, chills, rigors, pruritus, hemodynamic changes) the following orders will be followed and physician will be notified:<br>- Mild reaction: Pause infusion for 10 minutes, resume infusion at a minimum 50% reduction in rate after symptoms have resolved.<br>- Moderate reaction: Pause infusion, administer Diphenhydramine 25 mg IV; administer Sodium Chloride 0.9% 500mL IV bolus.<br>If symptoms persist, administer remaining Diphenhydramine 25 mg IV. Administer Diphenhydramine IM if no IV access. Notify Pharmacist.<br>- Severe reaction (w/breathing problems): CALL 911, administer Epinephrine 0.3 mg IM; administer Diphenhydramine 50mg IV x 1 dose; administer Sodium Chloride 0.9% 500mL IV bolus. Administer Diphenhydramine IM if no IV access. |                                                                                                                             |

## Prescriber Information

|                                                       |           |                |                                                    |
|-------------------------------------------------------|-----------|----------------|----------------------------------------------------|
| Physician Name                                        |           | Office Contact |                                                    |
| Practice Address                                      |           | Practice Phone |                                                    |
| NPI#                                                  | License # | DEA#           |                                                    |
| Physician Signature Required - Substitution Permitted |           | Date           | Physician Signature Required - Dispense as Written |
|                                                       |           |                | Date                                               |