Ocrelizumab and hyaluronidase-ocsq (Ocrevus Zunovo) Patient Referral Form Patient Demographics Information SSN# DOB Patient Name Patient Address Primary Phone Cellular Phone Emergency Contact Name, Relationship **Emergency Contact Phone Number** Additional Documentation Needed Copy of insurance cards History and Physical Recent vitals including blood pressure - Screening for HBV infection Patient face sheet w/demographics - CBC w/diff, BMP, & CMP Patient Insurance Information Insurance Plan #1 Insurance Plan #2 Subscriber Name DOB Subscriber Name Policy Number Group ID Policy Number Group ID Patient Clinical Information Gender Height (inches) Weight (lbs.) Allergies (food/drug) \square M Statement of Medical Necessity / Primary Diagnosis Description of diagnosis: Medication Information / Prescription and Orders Medication: **Directions:** Quantity/Refills Dispense: #1 dose Ocrevus Zunovo Infuse SubQ per manufacturer guidelines over approximately 10 minutes. + refills ☐ Other Dosing: Date of Last Dose: ☐ MS dosing: 920 mg/23,00 units SubQ every 6 months **Premedications** Quantity/Refills Dispense: quantity #QS Give premedication(s) 30 minutes prior to infusion PRN refills (unless otherwise noted) Acetaminophen □ 325-650 mg PO OR □ _____ mg PO Dexamethasone □ 20 mg PO ☐ Other Diphenhydramine □ 25-50 mg PO Other **Adverse Reaction Orders** · Mild reaction: Pause infusion and resume after symptoms have resolved Moderate reaction: Pause infusion and give diphenhydrAMINE 50mg PO. Resume infusion after symptoms have resolved. Notify Pharmacist. Severe reaction (w/breathing problems): stop infusion, call 911 and administer EPHINEPHrine 0.3 mg or 0.15 mg IM (as determined by patient weight). Prescriber Information Physician Name Office Contact Practice Address License # Practice Phone Physician Signature Required - Substitution Permitted Date Physician Signature Required - Dispense as Written Date