Pediatric Subcutaneous Immune Globulin Patient Referral Form Patient Demographics Information SSN# DOB Patient Name Patient Address Primary Phone Cellular Phone Work Phone Emergency Contact Name, Relationship **Emergency Contact Phone Number** Additional Documentation Needed Copy of insurance cards Labs to include IgA level (within 1 year) - For immune deficiency, detailed infection Patient face sheet w/demographics - CBC w/dif, BMP, & CMP history, baseline IgG levels, vaccine responses History and Physical - Recent vitals including blood pressure - Blood type Patient Insurance Information Insurance Plan #1 Insurance Plan #2 DOB DOB Subscriber Name Subscriber Name Policy Number Group ID **Policy Number** Group ID **Patient Clinical Information** Height (inches) Gender Weight (lbs.) Allergies (food/drug) \square M \Box F Statement of Medical Necessity / Primary Diagnosis Neurology Referrals Immunology Referrals Description of diagnosis Description of diagnosis Medication Information / Prescription and Orders Medication Dose Directions Quantity / Refills ☐ Preferred Product Loading: ___ gms OR gm/kg Infuse SQ per manufacturer guidelines OR Dispense: given over _____ days hours. 1 months supply, refill x12mos gms OR gm/kg Titration rate according to pharmacy unless otherwise noted ☐ Other (rounded to the nearest vial size) protocol. ☐ No Preference SQ every week(s) First Dose? If NO, List Product Date of last Infusion Next Dose Due \square Y \square N Quantity / Refills Premedications ☐ Give premedication 30 minutes prior to infusion (generics will be dispensed) Dispense: mg po (Diphenhydramine 12.5 mg/5mL Liquid - #1 bottle) Quantity #QS + PRN refills Diphenhydramine: Acetaminophen: _ mg po (Acetaminophen 160 mg/5 mL oral susp - #1 bottle) unless otherwise noted ☐ Other Other: (Physician to specify) 🗆 EMLA topical cream (Lidocaine 2.5% and Prilocaine 2.5%): apply to SQ needle site prior to access PRN for pain upon needle insertion. ☑ Instruct patient or caregiver to hydrate pre/post infusion. ☐ Other (Physician to specify): Adverse Reaction Orders In the event of an infusion reaction (ie: fever, chills, backache, headache, rigors) the following orders will be followed and physician will be notified: Mild to moderate reaction: Diphenhydramine 1 mg/kg (≤ 25 kg) or 25 mg (> 25 kg) PO x1 dose and pause infusion until symptoms resolve. May repeat every 6 to 8 hours. Maximum 50 mg/dose. Dispense Diphenhydramine 12.5 mg/5 mL liquid. Severe reaction (w/breathing problems): CALL 911 and administer Epinephrine IM 0.1 mg, 0.15 mg, or 0.3 mg (as determined by patient wt). Prescriber Information Physician Name Office Contact Practice Address Practice Phone License # Physician Signature Required - Substitution Permitted Physician Signature Required - Dispense as Written Date Date